

Summary: Climate Change in Tuvalu 2022


Historical and Recent Variability, Extremes and Change



COSPPac
Climate and Oceans Support
Program in the Pacific

Figure 2:

Annual longest run of consecutive dry days (bar graph) and maximum daily rainfall (line graph) at Funafuti. Straight lines indicate linear trends for dry days (in black) and maximum daily rainfall (in blue). Diamonds indicate years with insufficient data for one or both variables.



There has been little long-term change in meteorological drought over time.

Air Temperature has increased

Average annual temperatures at Funafuti have increased by 0.19 °C per decade since 1951. Dry season (May–October) temperatures and night-time temperatures warmed faster than wet season (November–April) and daytime temperatures.

The number of hot days and warm nights has increased, and the number of cool days and cold nights has decreased at Funafuti. Since 1951, the number of hot days has increased by 29 days per decade. Hot days have a maximum temperature above 31.1–32.4 °C, depending on the time of year.

The number of days where air conditioning is required to cool a building down to 25 °C has increased by 71 days per decade, indicating that energy demand for cooling has increased significantly since 1951.

Long-term increases in both average temperature and temperature extremes in the Pacific are likely driven by human-associated climate change, due to the rate of the observed changes and consistency with global trends that have been attributed to climate change (PCCM, 2021).



Tropical cyclone severity has decreased


In the greater Southwest Pacific, the total number of **severe** tropical cyclones¹ has decreased over the last 40 seasons. There has been little change in the total number of tropical cyclones of any category in the southwest Pacific. The number of tropical cyclones that became severe events has marginally declined.

Tropical cyclones usually affect Tuvalu during the southern hemisphere tropical cyclone season, which is from November to April, but also occasionally occur outside the tropical cyclone season.

The number of tropical cyclones occurring in Tuvalu's Exclusive Economic Zone (EEZ) varies considerably from one year to the next (Figure 3). Tropical cyclones were most frequent in El Niño years (12 cyclones per decade), followed by neutral years (10 cyclones per decade) and least frequent in La Niña years (3 cyclones per decade).

Figure 3:

Number of tropical cyclones passing within the Tuvalu EEZ per season. Each season is defined by the ENSO status, with light blue being an El Niño year, dark blue a La Niña year and grey showing a neutral ENSO year. The 11-year moving average is presented as a purple line and considers all years.



Due to this high interannual variability and the relatively small number of tropical cyclones passing through any country's EEZ since reliable records began, individual country analysis of long-term trends in frequency and intensity is not possible.


¹ A 'severe' tropical cyclone is defined as having a minimal central pressure of <970 hectopascals (hPa). Pressure is often used when comparing intensity of tropical cyclones.

Sea surface temperature has increased

Sea surface temperatures averaged across Tuvalu's EEZ increased by 0.22 °C per decade since 1982 (Figure 4).

Figure 4:

Sea surface temperature from satellite observations averaged across the Tuvalu EEZ, shown as the orange line. The blue line shows the linear regression trend.



Globally, sea surface temperature is one of the most widely used indicators used to monitor human-associated climate change. Modes of climate variability influence sea surface temperatures on an interannual and decadal/multi-decadal basis; however, climate change is a driver of the long-term positive trend (PCCM, 2021).

Sea surface temperatures at Funafuti tend to be warmest in November/December and then again around April/May, reaching, on average, a maximum of almost 30 °C. Sea surface temperatures are coolest in August, reaching, on average, a minimum of 29 °C. Hourly temperatures can be up to 2 °C higher or lower than these monthly averages at Funafuti and may differ at other locations in Tuvalu.




Sea level has increased

A combination of sea level rise and land subsidence has increased sea level at the Funafuti tide gauge by 4.5 mm per year since 1993 (Figure 5). The number of hours per month that sea level has exceeded the 99th percentile of historical maximum sea level has increased since 2010. Peak sea levels typically occur between January and April.

Figure 5:

The effect of sea level rise and land subsidence on local sea level.



The long-term trend in sea level from satellite altimetry across Tuvalu's EEZ is 3.5–4.5 mm per year since 1993. This trend is higher than the global average trend.



The rise in Pacific mean sea level since 1993 is primarily attributable to global warming. Naturally-occurring modes of climate variability in the Pacific region - for example, the El Niño–Southern Oscillation (ENSO) on interannual time scales, and the IPO (Interdecadal Pacific Oscillation)/PDO (Pacific Decadal Oscillation) on decadal to multi-decadal time scales - influence sea level and can amplify or dampen the underlying trends arising from global warming (PCCM, 2021)

Local flooding, Tuvalu



Waves

Waves at Funafuti come from the northeast to the southeast. On average, Funafuti experiences approximately three extreme wave events – defined as reaching or exceeding a wave height of 2.26 m per year.

There has been no long-term change in average annual wave height since 1979. Wave height, wave period (the time interval between two waves) and wave direction changes from month to month with the seasons and, to a lesser degree, year to year with climate variability modes. The highest waves usually occur between June to September.



Further reading

For more information, refer to Climate Change in the Pacific 2022: Historical and Recent Variability, Extremes and Change. Climate and Oceans Support Program in the Pacific. Fifteen country chapters are available at <https://purl.org/spc/digilib/doc/kskiv>

For more information on Pacific-wide observed and future trends in climate indicators, see the Pacific Islands Climate Change Monitor 2021, available at https://www.pacificmet.net/sites/default/files/inline-files/documents/PICC%20Monitor_2021_FINALpp_0.pdf

Historical climate trends and basic climate information from observation sites across the Pacific Islands are available through the web-based Pacific Climate Change Data Portal at www.bom.gov.au/climate/pccsp

Information about future climate change can be found in the 'NextGen' Projections for the Western Tropical Pacific country reports <https://www.csiro.au/en/research/environmental-impacts/climate-change/pacific-climate-change-info>





Local fisherman in Tuvalu

The content of this brochure is an outcome of the high degree of cooperation and collaboration that exists between the implementing partners of the Australian Aid funded Climate and Oceans Support Program in the Pacific (COSPPac), specifically the Bureau of Meteorology (the Bureau), the Pacific Community (SPC) and Pacific Regional Environmental Programme (SPREP), together with the valuable ongoing support from the national meteorological services in the 15 partner countries and territories. Publication support has been provided through New Zealand Aid Programme.



For more detailed information on the climate of Tuvalu and the Pacific, see: *McGree, S., G. Smith, E. Chandler, N. Herold, Z. Begg, Y. Kuleshov, P. Malsale and M. Ritman. 2022. Climate Change in the Pacific 2022: Historical and Recent Variability, Extremes and Change. Climate and Oceans Support Program in the Pacific. Pacific Community, Suva, Fiji.*



Contact the Tuvalu Meteorological Service:

web: <https://tuvmet.tv/>

Facebook: @Tuvalu Meteorological Service

Twitter: @TuvaluMET

phone: +688 20736