

A biometric study to determine the economic and nutritional value of sea cucumbers (Holothuroidea: Echinodermata) collected from Algeria's shallow water areas

Asmaa Mecheta¹ and Karim Mezali¹

Abstract

Traditionally in Algeria, sea cucumbers were exploited at a very small scale. Nowadays, however, sea cucumber fisheries are emerging in the Mediterranean Sea due to the high commercial value of these organisms. Three economically important species – *Holothuria (Roweothuria) poli*, *H. (Holothuria) tubulosa* and *H. (Roweothuria) arguinensis* – are common in shallow water areas along Algeria's coast. In order to enhance and highlight the economic and nutritional value of these unexploited holothurian species, a biometric study estimating the amount of water and pepsin digestion of the dry body wall, was undertaken. The three holothurians species were collected at two stations: Stidia and Hadjadj near the port city of Mostaganem. After their transformation into dried products, called beche-de-mer, we noticed a considerable decrease in their length (50–73%) and weight (88–93%). The dried body wall analysis showed a high moisture rate (over 80%) and pepsin digestibility (varying between 25% and 52%). We compared our results with those obtained for other holothurian species commonly exploited in the Indo-Pacific region.

Keywords: holothurians, biometrics, dried product, digestibility, Algerian west coast

Introduction

The demand for beche-de-mer by consumer countries is constantly rising, resulting in increasingly intensive fisheries and a considerable decrease, and in some areas disappearance, of natural populations (Samyn et al. 2006). For that reason, Asian markets are in search of new target species, mainly from the Mediterranean Sea and the northeast Atlantic Ocean (Gonzalez-Wangüemert et al. 2015, 2016). In these regions, the most important species are *Holothuria (Roweothuria) poli* Delle Chiaje 1823, *Holothuria (Holothuria) tubulosa* Gmelin 1791, *Holothuria (Platyperona) sanctori* Delle Chiaje 1823, *Holothuria (Panningothuria) forskali* Delle Chiaje 1823 and *Parastichopus regalis* Cuvier, 1817.

Recently, *Holothuria (Roweothuria) arguinensis* Koehler and Vaney 1906, an invasive species originating from the Atlantic Ocean, has been reported in the Mediterranean (González-Wangüemert and Borrero-Pérez 2012; Mezali and Thandar 2014). In Algeria, holothurians are exploited on a very small scale (mainly as fishing bait) and no attempt has been made to process these species into dried product. In order to enhance and highlight the economic

and the nutritional value of these unexploited holothurian species, a biometric study estimating the amount of water and pepsin digestion of the dried body wall was conducted.

Methodology

Sampling was carried out during the summer of 2018. Three batches of 30 individuals of each species – *H. poli*, *H. tubulosa* and *H. arguinensis* – were harvested from two sampling sites on the Algerian west coast: Stidia (35° 50'N, 0° 00'E) and Hadjadj (36° 06'N, 0° 20'E) (Fig. 1). Each individual from each species was isolated in a plastic bag for later analysis.

Biometry

In the laboratory, each sea cucumber individual was cleaned and then measured. The contracted length was obtained according to the standardised method described by Mezali (1998) using a semi-cylindrical scale (± 0.5 mm). Each measured individual was kept straight, and then compressed by hand until the coelomic fluid was expelled. After that, the individuals were dissected by making an incision

¹ Protection, Valuation of Marine and Coastal Resources and Molecular Systematics Laboratory, Department of Marine Sciences and aquaculture, Faculty of Natural Sciences and Life, Abdelhamid Ibn Badis University-Mostaganem, 27000, PO Box 227, Algeria

* Author for correspondence: asmaamecheta@gmail.com




Figure 1. Detailed map of the two sampling sites (A) (black crosses).

of the trivium using a scalpel blade (Purcell 2014). Each individual was eviscerated, its gut and viscera removed, rinsed with tap water, drained and finally weighed: wet weight of the body wall (WW) ($\pm 0.001\text{g}$). The wet weight included the body wall, longitudinal muscle and the buccal bulb (Kazanidis et al. 2010; Mezali et al. 2014), and gave more accurate information than total body weight, which varies considerably, depending on the amount of water in the respiratory trees and the gut contents (mainly composed by exogenous materials) (Mezali 1998). The trivium thickness was measured using a caliper ($\pm 0.1\text{ mm}$). The moisture content of the body wall was measured according to the Official Methods of Analysis (AOAC 950.01 1990). The results have been expressed in percentages (%) (the mean of three replicates). The dried body wall of each individual was placed into a labelled plastic bag and stored in a freezer ($-4\text{ }^{\circ}\text{C}$) for latter biochemical analysis. Fresh sea cucumbers were processed into dried products based on the simple, traditional methods described by Purcell (2014), and organised in three basic steps: cooking, salting and sun drying. Freshly collected holothurians were placed in a large container and boiled in sea water ($60\text{--}80\text{ }^{\circ}\text{C}$), and afterwards were placed in cold water ($15\text{ }^{\circ}\text{C}$ for 1–2 h). An incision (with a scalpel blade) was made along the trivium, starting at 3 cm from each end, and the samples then gutted and rinsed in tap water. The specimens were kept in bins and entirely covered with salt for 2–5 days. Sea cucumbers were boiled again in sea water for $\frac{1}{2}$ h, until a hard and chewy consistency was obtained. Finally, they were sun dried for 4–14 days.

In vitro digestion of the dried body wall with pepsin

Dried body wall of *H. poli*, *H. tubulosa* and *H. arguinensis* were digested *in vitro* according to the method

of Escudero et al. (2010) and amended by Wen et al. (2015). For this, 0.5 g of dried body wall of each species was weighed ($\pm 0.001\text{ g}$) (Jinghua JA3003, Shanghai, China) and then homogenised twice in 2 mL of distilled water for 30 s at $9,500\text{ rpm min}^{-1}$ and then twice again for 30 s at $13,500\text{ rpm min}^{-1}$ with a 30 s cooling time between the two centrifugations. The homogenates were adjusted to a pH of 2 with an acid (HCl 1M). Pepsin was added in a 1:31.25 ratio based on the body wall mass. The mixture was maintained at $37\text{ }^{\circ}\text{C}$ for 2 h, and then was made inactivate by adjusting the pH to 7.5 using a base (NaOH 1M). The undigested body wall was dried in an oven at $60\text{ }^{\circ}\text{C}$ for 72 h and then weighed. The *in vitro* digestibility was evaluated according to the following formula:

$$\text{DT} = (1 - \text{Wi} / \text{Wt}) \times 100\%$$

Where DT = digestibility (%);

Wi = dried weight of body wall after digestion;

Wt = total weight of the body wall of the dried product before digestion.

Results and discussion

Biometry

Our specimens demonstrated larger class sizes than those recorded by Mezali (1998) and Purcell et al. (2009) (Table 1). This difference could be due to several factors, such as a rapid growth due to a decrease in predation (Olaya-Restrepo et al. 2018). Although few predators feed on holothurians, several anti-predation mechanisms have been described in the literature for sea cucumbers (Francour 1997), and some of these mechanisms appear to be related to the animal's size. Moreover, this could also be linked to the high productivity of

Posidonia oceanica meadows, the preferred habitat for *H. poli* and *H. tubulosa* (Mezali 2004). In addition, the smaller the sediment particles, the higher the quantity of organic matter that can be ingested by holothurians, due to a higher volume-to-surface ratio that has previously been demonstrated to promote the adhesion of organic matter (Mezali and Soualili 2013). Moreover, Plotieau et al. (2013) suggested that the nutritional value of fine sediment would be higher than that for coarse sediments due to the greater number of nutritious microorganisms.

It is still, however, important to point out that estimating total body length is not a reliable measure. In fact, the error is likely to be higher than the actual weight because it is very difficult to obtain complete relaxation of holothurians' bodies; the length indeed changes significantly by contraction and elongation of their body (Battaglione et al. 1999). Therefore, weight measurements are more reliable (Mezali 1998). Based on this assertion, the average eviscerated weight was measured for the three species, and the maximum values were recorded as: *H. tubulosa* (112.64 ± 45.57 g), *H. arguinensis* (244.8

± 61.26 g) and *H. poli* (90.22 ± 16.07 g) (Table.1). The relationship between length and weight differ between species, depending on the shape and strength of the body (Cone 1989). This is also due to food availability, species biology, growth rates and fishing pressure (González-Wangüemert et al. 2018).

A difference in body wall thickness could be observed between the three studied species: (5.99 ± 0.8 mm) in *H. tubulosa*, (3.44 ± 0.56 mm) in *H. poli* and (4.53 ± 1.03 mm) in *H. arguinensis* (Table1). According to González-Wangüemert et al. (2018), food is used differently by the two species – *H. poli* and *H. tubulosa* – with *H. poli* increasing in length while *H. tubulosa* increasing its body wall robustness (thickness of the body wall) and weight. Our results are not in agreement with this hypothesis because *H. tubulosa* presents the largest size classes, and a weight and body wall thickness greater than those recorded for *H. poli*, despite its more rigid body wall. Regarding *H. arguinensis*, although its specific growth model has already been determined by Olaya-Restrepo et al. (2018), there is no information on the relationship

Table 1. Biometric measurements obtained for the three studied holothurians species compared with other species of the Mediterranean and Indo-Pacific regions (n = sample size = 30). CL = contracted length (mm); T = thickness of the body wall (mm); WW = wet weight of the body wall (g); DW = dry weight of the body wall (g); and DL = dry length of the body wall (mm).

Species	CL (mm)	T (mm)	WW (g)	DW (g)	DL (mm)	WW/DW	References
<i>Actinopyga echinites</i>	190.00 ± 3.00	-	231.00 ± 14.00	35.00 ± 2.00	80.00 ± 2.00	-	Purcell et al. (2009)b
<i>A. palauensis</i>	270.00 ± 7.00	-	985.00 ± 44.00	165.00 ± 11.00	150.00 ± 5.00	-	Purcell et al. (2009)b
<i>A. spinea</i>	270.00 ± 10.00	-	735.00 ± 39.00	99.00 ± 11.00	130.00 ± 10.00	-	Purcell et al. (2009)b
<i>Holothuria arguinensis</i>	261.00 ± 55.16	4.53 ± 1.03	244.80 ± 61.26	15.51 ± 3.60	67.30 ± 5.22	15.78	this study
<i>H. forskali</i>	108.04 ± 28.90	-	-	-	-	8.89	Mezali (1998)
<i>H. lessoni</i>	310.00 ± 10.00	-	1456.00 ± 50.00	221.00 ± 7.00	280.00 ± 10.00	-	Purcell et al. (2009)b
<i>H. poli</i>	144.03 ± 23.07	3.44 ± 0.56	90.22 ± 16.07	4 12.85 ± .59	71.60 ± 8.68	7.02	this study
<i>H. poli</i>	91.03 ± 43.10	-	-	-	-	6.44	Mezali (1998)
<i>H. sanctori</i>	115.60 ± 27.90	-	-	-	-	-	Mezali (1998)
<i>H. sanctori</i>	225 ± 38.90	02.69 ± 1.95	76.90 ± 18.37	-	-	-	Mezali et al. (2014)a
<i>H. tubulosa</i>	187.30 ± 39.79	5.96 ± 1.26	112.64 ± 45.57	12.95 ± 1.32	74.20 ± 9.80	8.69	this study
<i>H. tubulosa</i>	120.15 ± 38.52	-	-	-	-	6.53	Mezali (1998)
<i>H. whitmaei</i>	250.00 ± 8.00	-	1174.00 ± 45.00	213.00 ± 14.00	150.00 ± 4.00	-	Purcell et al. (2009)b

^a anesthetised total length

^b length measurement criteria not indicated

between food availability and the strength of the body wall of this species. After being processed into beche-de-mer, the dried sea cucumbers were weighed and measured to estimate the weight and length loss. The results demonstrate a significant decrease in weight and size in all studied species (Table 1, Fig. 2). Lengths varied between 40.10% and 74.21%, and weights ranged from 85.72% to 93.65%. According to Conand (1990), after the drying process, the weight decreases between 90% and 97%, depending on the species. The salting phase allows dehydration of the sea cucumbers and removes part of the water contained in the body wall prior to sun drying. Similar studies conducted by Purcell et al. (2009) and Lavitra et al. (2009) on sea cucumber species exploited in New Caledonia and southwestern Madagascar, confirm our results as these authors have also found considerable decreases in the length and weight of fresh and dried sea cucumbers.

The ratio between wet weight and dry weight was calculated for all species. Considerable variations were observed between the three studied species (Table 1). The wet weight to dry weight ratio for both *H. poli* and *H. tubulosa* approximates the standard value 10:1 established for sea cucumbers by Newell and Courtney (in Astall and Johns 1991). But the ratio obtained for *H. arguinensis* is greater than the standard value 10:1. Our results also differ from those obtained by Mezali (1998) for similar species: *H. poli* (6.44:1) and *H. tubulosa* (6.53:1).

Water content and *in vitro* digestion of the dry integument

Our results show very high moisture content for the three studied holothurian species (Table 2). Sea cucumbers generally have a higher moisture content than that of fish and seafood (Chang-Lee et al. 1989).

Table 3 shows the digestibility rate of the body wall of the three studied sea cucumbers by the digestive enzyme pepsin. The digestibility




Figure 2. Dried sea cucumbers A. *H. poli*; B. *H. tubulosa*; C. *H. arguinensis* after processing into beche-de-mer, according to the traditional method. (image: Asmaa Mecheta)

Table 2. Water content (%) in the body wall of the three studied holothurians compared with Indo-Pacific species (no. replicates = 3).

Species	Body wall water content (%)	References
<i>Actinopyga mauritiana</i>	76.54 ± 0.09	Haider et al. (2015)
<i>Holothuria arenicola</i>	72.12 ± 0.25	Haider et al. (2015)
<i>H. arguinensis</i>	88.13 ± 4.16	this study
<i>H. fuscogilva</i>	84.34 ± 0.72	Fawzya et al. (2015)
<i>H. leucospilota</i>	81.41 ± 0.60	Omran (2013)
<i>H. poli</i>	93.53 ± 1.95	this study
<i>H. poli</i>	81.24 ± 0.40	Aydin (2009)
<i>H. scabra</i>	85.76 ± 0.30	Omran (2013)
<i>H. tubulosa</i>	81.99 ± 1.97	this study
<i>H. tubulosa</i>	84.30 ± 0.20	Aydin (2009)
<i>Parastichopus californicus</i>	04.03 ± 0.19	Bechtel et al. (2013)
<i>Stichopus herrmanni</i>	10.20 ± 0.32	Wen et al. (2010)
<i>Thelenota ananas</i>	90.81 ± 2.08	Fawzya et al. (2015)
<i>T. anax</i>	01.20 ± 0.06	Wen et al. (2010)

Table 3. Digestibility (%) of the holothurian body wall when digested using pepsin.

	Digestibility (%)	References
<i>Holothuria arguinensis</i>	53.56 ± 3.41	this study
<i>H. poli</i>	34.68 ± 8.66	this study
<i>H. tubulosa</i>	25.96 ± 2.04	this study
Pork	47.22	Wen and al. (2015)
Fish	46.98	Wen and al. (2015)
Chicken	44.67	Wen and al. (2015)
Beef	42.75	Wen and al. (2015)

of *H. arguinensis* ($53.56 \pm 3.41\%$) was significantly higher than that of *H. poli* ($34.68 \pm 8.66\%$) and *H. tubulosa* ($25.96 \pm 2.04\%$) (Table 3). The digestibility level of *H. arguinensis* was more or less similar to that obtained by Wen et al. (2015) for commonly consumed meats (pork, fish, chicken and beef). This digestibility variation was attributed to the changes in protein bioavailability (depending on the protein decomposition rate when degraded by digestive enzymes), peptide size (Hur et al. 2011; Pennings et al. 2013), and protein and collagen contents that constitute the major components of the holothurians body wall (about 70% of the total body wall protein) (Saito et al. 2002; Wen et al. 2015). This latter factor may be significant due to the presence of glycosaminoglycans, which are widely distributed among the bundles of collagen fibers (Kariya et al. 1990).

Conclusion

This biometric study of three aspidochirotid holothurian species – *H. poli*, *H. tubulosa* and *H. arguinensis* – before and after processing into beches-de-mer showed that our specimens are smaller in size and weight than those exploited in the Indo-Pacific region. A lower trophic factor combined with less organic matter content mostly explains these outcomes. Despite these two factors, the studied holothurians species showed significant nutritional value as their body wall is degraded by the digestive enzyme pepsin. Thus, they could become common sources of protein and other nutrients.

Acknowledgments

The first author thanks the microbiology and animal biology laboratory teams of the Natural Sciences and Life Faculty, Abdelhamid Ibn Badis University at Mostaganem, for their help in carrying out the experiments. Also, thanks to scuba diver Mr Abdelkader who helped during the field work.

References

- Astall C.M. and Johns M.B. 1991. Respiration and biometry in the sea cucumber *Holothuria forskali*. *Journal of the Marine Biological Association of the United Kingdom* 71:73–81.
- Aydin M., Huseyin S., Bekir T., Yilmaz E., Sevim K. 2010. Proximate composition and fatty acid profile of three different fresh and dried sea cucumber commercial from Turkey. *International Journal of Food Science and Technology* 46: 500–508.
- Battaglione S.C., Seymour J.E. and Ramofafia C. 1999. Survival and growth of cultured juvenile sea cucumbers, *Holothuria scabra*. *Aquaculture* 178:293–322.
- Bechtel P.J., Oliveira A.C.M., Demir N., Smiley S. 2013. Chemical composition of the giant red sea cucumber *Parastichopus californicus*, commercially harvested in Alaska. *Food Science & Nutrition* 1:63–73.
- Chang-Lee M.V., Price R.J. and Lampila L.E. 1989. Effect of processing on proximate composition and mineral content of sea cucumbers (*Parastichopus* spp.). *Journal of Food Science* 54:567–572.
- Conand C. 1990. The fishery resources of Pacific Island countries. Part 2: Holothurians. *FAO Fisheries Technical Paper*, No. 272.2. Rome: Food and Agriculture Organization. 143 p.
- Cone R. S. 1989. The need to reconsider the use of condition indices in fishery science. *Transactions Journal of the American Fisheries Society* 118:510–514.
- Cuvier G. 1817. Les Annélides. p. 515–532. In : *Le règne animal distribué d'après son organisation, pour servir de base à l'histoire naturelle des animaux et d'introduction à l'anatomie comparée*. Volume 2. Contenant les Reptiles, les Poissons, les Mollusques et les Annélides. Paris, France : Deterville.
- Della Chiaje S. 1823. Descrizione zoologica ed anatomica di alcune specie di Oloturie. p. 77–116. In: *Memorie su la storia e notomia degli animali senza vertebre del Regno di Napoli*. Volume 1 Fratelli Fernandes, Napoli. pls. 6–8.
- Escudero E., Sentandreu M.A. and Toldrá F. 2010. Characterization of peptides released by in vitro digestion of pork meat. *Journal of Agricultural and Food Chemistry* 58:5160–5165.
- Fawzya Y.N., Januar H.I., Susilowati, R., Chasanah E. 2015. Chemical composition and fatty acid profile of some Indonesian sea cucumbers. *Squalen Bulletin of Marine and Fisheries Postharvest Biotechnology* 10:27–34.
- Francour P. 1997. Predation on holothurians: A literature review. *Invertebrate Biology* 116:52–60.
- Gmelin J.F. 1791. Vermes. p. 3021–3910. In: Gmelin J.F. (ed). *Caroli a Linnaei Systema Naturae per Regna Tria Naturae*, ed. 13. Tome 1(6). Lipsiae: G.E. Beer.
- González-Wangüemert M. and Borrero-Pérez G. 2012. A new record of *Holothuria arguinensis* colonizing the Mediterranean Sea. *Marine Biodiversity Records* 5:1–4.
- González-Wangüemert M., Valente S. and Aydin M. 2015. Effects of fishery protection on biometry and genetic structure of two target sea cucumber species from the Mediterranean Sea. *Hydrobiologia* 743:65–74
- Gonzalez-Wangüemert M., Domínguez-Godino J.A. and Canovas F. 2018. The fast development of sea cucumber fisheries in the Mediterranean and NE Atlantic waters: From a new marine resource to its over-exploitation. *Ocean and Coastal Management* 151:165–177.

- González-Wangüemert M., Valente S., Henriques F., Domínguez-Godino J.A. and Serrão E.A. 2016. Setting preliminary biometric baselines for new target sea cucumbers species of the NE Atlantic and Mediterranean fisheries. *Fisheries Research* 179:57–66.
- Haider M.S, Sultana R., Jamil Lakht-e-Zehra K., Tarar O.M., Shirin K., Afzal W. 2015. A Study on proximate Composition Amino Acid Profile, Fatty Acid Profile and Some Mineral Contents in Two Species of Sea Cucumber. *Journal of Animal and Plant Sciences* 25:168–175.
- Hur S.J., Lim B.O., Decker E.A. and Mc Clements D.J. 2011. In vitro human digestion models for food applications. *Food Chemistry* 125:1–12.
- Kariya Y., Watabe S., Ochiai Y., Murata K. and Hashimoto K. 1990. Glycosaminoglycan involved in the cation-induced change of body wall structure of sea cucumber *Stichopus japonicus*. *Connective Tissue Research* 25:149–159.
- Kazanidis G., Antoniadou C., Lolas A.P., Neofitou N., Vafidis D., Chintiroglou C. and Neofitou C. 2010. Population dynamics and reproduction of *Holothuria tubulosa* (Holothuroidea: Echinodermata) in the Aegean Sea. *Journal of the Marine Biological Association of the United Kingdom* 90:895–901.
- Lavitra T., Rachele D., Rasolofonirina R., Jangoux M. and Eeckhaut I. 2009. Processing and marketing of holothurians in the Toliara region, southwestern Madagascar. *SPC Beche-de-mer Information Bulletin* 28:24–33.
- Mezali K. 1998. Contribution à la systématique, la biologie, l'écologie et à la dynamique des populations de cinq espèces d'holothuries aspidochirotes (*Holothuria tubulosa*, *Holothuria poli*, *Holothuria stellati*, *Holothuria forskali* et *Holothuria sanctori*) de l'herbier à *Posidonia oceanica* (L.) Delile de la presqu'île de Sidi-Fredj- Algérie (Thèse de Doctorat d'état. Alger, Algérie : USTHB) 208. 8 p.
- Mezali K. 2004. Micro-répartition des holothuries aspidochirotes au sein de l'herbier de *Posidonies* de la presqu'île de Sidi-Fredj (Algérie). p. 534. CIESM (ed.). Rapport du 37^e Congrès de la Commission internationale pour l'exploration scientifique de la Mer Méditerranée. Monaco : Commission internationale pour l'exploration scientifique de la Mer Méditerranée. 574 p.
- Mezali K. and Soualili D.L. 2013. The ability of holothurians to select sediment particles and organic matter. *SPC Beche-de-mer Information Bulletin* 33:38–43.
- Mezali K. and Thandar A. 2014. First record of *Holothuria (Roweothuria) arguinensis* (Echinodermata: Holothuroidea: Aspidochirotida: Holothuriidae) from the Algerian coastal waters. *Marine Biodiversity Records* 4:7–40.
- Mezali K., Soualili D.L., Neghli L. and Conand C. 2014. Reproductive cycle of the sea cucumber *Holothuria (Platyperona) sanctori* (Holothuroidea: Echinodermata) in the southwestern Mediterranean Sea: interpopulation variability. *Invertebrate Reproduction and Development* 58(3):179–189.
- Olaya-Restrepo J., Erzini K. and González-Wangüemert M. 2018. Estimation of growth parameters for the exploited sea cucumber *H. arguinensis* from South Portugal. *Fishery Bulletin* 116:1–8.
- Omran N. 2013. Nutritional value of Some Egyptian sea cucumbers. *African Journal of Biotechnology* 12:35.
- Pennings B., Groen B.B., van Dijk J.W. and de Lange A. 2013. Minced beef is more rapidly digested and absorbed than beef steak, resulting in greater postprandial protein retention in older men. *The American Journal of Clinical Nutrition* 98:121–128.
- Plotieau T., Baele J.M., Vaucher R., Hasler C.A., Koudad D. and Eeckhaut I. 2013. Analysis of the impact of *Holothuria scabra* intensive farming on sediment. *Cahiers de Biologie Marine* 54:703–711.
- Purcell S.W. 2014. Processing sea cucumbers into beche-de-mer: A manual for Pacific Island fishers. Noumea, New Caledonia: Southern Cross University, Lismore, and the Secretariat of the Pacific Community. 44 p.
- Purcell S.W., Gossuin H. and Agudo N.S. 2009. Changes in weight and length of sea cucumbers during conversion to processed bêche-de-mer: Filling gaps for some exploited tropical species. *SPC Beche-de-mer Information Bulletin* 29:3–6.
- Saito M., Kunisaki N., Urano N. and Kimura S. 2002. Collagen as the major edible component of sea cucumber (*Stichopus japonicus*). *Journal of Food Science* 67:1319–1322.
- Samyn Y., Van den Spiegel D. and Massin C. 2006. Taxonomie des holothuries des Comores. *AbcTaxa* vol 1 : i-iii. 130 p.
- Wen S., Zhou G., Song S., Xu X., Voglmeir J., Liu L, Zhao F., Li M., Li L., Yu X., Bai Y. and Li C. 2015. Discrimination of in vitro and in vivo digestion products of meat proteins from pork, beef, chicken, and fish. *Proteomics* 15:3688–3698.