

# Assessing rehydration protocols on dried sea cucumber *Holothuria arguinensis*

Mercedes González-Wangüemert<sup>1\*</sup> and Jorge A. Domínguez-Godinoa<sup>1</sup>

## Abstract

Sea cucumbers have been traditionally consumed as a tonic food in Asia for thousands of years; between 80% and 90% of sea cucumbers are dried to prevent their autolysis and to facilitate their storage, exportation and sale. Although in recent years, an increase of fresh and frozen sea cucumber products has been observed in the Asiatic markets.

Scarce information about rehydration protocols has been published on sea cucumbers until now. *Holothuria arguinensis* is one of the new target species caught from NE Atlantic. In this paper, its rehydration ratio (Rr) is assessed using two different protocols and compared with ratios obtained with other commercial species such as *Holothuria scabra* and *Apostichopus japonicus*. The highest value of rehydration rate on *H. arguinensis* (Rr = 3.78) was relatively close to the value registered for *Apostichopus japonicus* (Rr = 4.07) by other authors, and higher than the value obtained for *H. scabra* (Rr = 3.05). The obtained results are very valuable to improve the rehydration protocols used on *H. arguinensis*, allowing the maximum recovery rates of weight and length from the dried product.

## Introduction

*Holothuria (Roweothuria) arguinensis* Koehler and Vaney, 1906, is being considered as a target species in the expanding sea cucumber fishery (González-Wangüemert et al. 2016, 2018). This species had been considered to be a north-eastern Atlantic species that is distributed from Portugal to Morocco and Mauritania, including the Canary Islands (González-Wangüemert and Borrero-Pérez 2012). It has not been found in other Macaronesian Islands such as Açores, Selvagens or Madeira, or in the Cape Verde Archipelago (Pereira 1997; Borrero-Pérez et al. 2010; Micael et al. 2012). However, its geographical distribution is changing, and includes the colonisation of the Mediterranean Sea (González-Wangüemert and Borrero-Pérez 2012; Mezali and Thadar 2014).

Sea cucumbers have been traditionally consumed as a tonic food in Asia and the Middle East for thousands of years (Zhang et al. 2016); they contain certain nutrients, which are beneficial to health. In the case of *H. arguinensis* high levels of protein were found and its lipid profile was rich in polyunsaturated fatty acids (PUFA) (Rogattz et al. 2016).

In recent years, sea cucumber consumption has been increasing in the Asian region with fisheries focusing on exportation of products in at least 70 countries (Purcell et al. 2012). More than 66

species are being fished, their dried product (the beche-de-mer) reaching prices up to USD 500 kg<sup>-1</sup> (Purcell et al. 2013). Sea cucumber-derived food products, including instant sea cucumber, sea cucumber capsules and sea cucumber caplets, can be found on the market (Zhang et al. 2016).

Usually, fresh sea cucumbers autolysis occurs rapidly after being harvested, due to the presence of autolytic enzymes in their body. Therefore, between 80% and 90% of sea cucumbers are dried to prevent autolysis and to facilitate their storage, exportation and sale (Aydin 2008; Zhang et al. 2016; González-Wangüemert et al. 2018). Although in recent years, an increase of fresh and frozen sea cucumber products has been observed in Asian markets (Purcell et al. 2014; Sze and Conand 2015).

Rehydration of dehydrated food is a complex phenomenon that is affected by numerous factors including medium characteristics, pre-soaking time, number of rehydration times, size of the food sample, and drying method (Geng et al. 2015). Most of the rehydration protocols of dried sea cucumbers are done in cold, clean water for 2–3 days before further processing or cooking (Duan et al. 2008).

The published information about sea cucumber rehydration process is scarce (Fukunaga et al. 2004; Liu and Ko 2002; Xiang et al. 2007). Hong et al. (2014)

<sup>1</sup> CCMAR- Centro do Ciências do Mar, Universidade do Algarve, Gambelas, 8005-139 Faro, Portugal

\* Author for correspondence: mwanguemert@ualg.pt

who worked with different rehydration methods on hot air-dried sea cucumber, found a rehydration ratio ranging from 5 to 6 for *Apostichopus japonicus*. Zhang et al. (2016) used ultrasound-assisted rehydration and obtained rates ranging from 2.71 to 4.35 for the same species. No information on the rehydration of *H. arguinensis*, a newly targeted species from the Mediterranean and NE Atlantic, has been published. Therefore, our objective has been to assess the rehydration ratio (Rr) of *H. arguinensis* using two different protocols and to compare it with ratios obtained with other commercial species, either by us (*Holothuria scabra*) or by other authors (*A. japonicus*).

## Material and methods

Two methodologies for rehydration of four *H. arguinensis* specimens from Ria Formosa (Southern Portugal) were used (Figure 1): A) the dried sea cucumbers were soaked in filtered water (filter of active carbon) during 96 hours at 4°C; the water was changed every 24 h; B) the dried sea cucumbers were submerged in filtered water during 3 hours, then boiled for 30 minutes, and finally, when the water temperature had decreased, they were conserved at 4°C overnight; this procedure was repeated at 40 h and 96 h. The recycling of water was carried out every 24 h, during the 96 h. One individual of *H. scabra* from Australia was also rehydrated using protocol-A.

The eviscerated weight ( $EW \pm 0.01$  g) and eviscerated length ( $EL \pm 0.1$  mm) for each individual were registered along the different phases of process. Water absorption by sea cucumbers was determined after rehydration. The rehydrated sea cucumbers were blotted with absorbent paper to remove excess water from their surface. Rehydrated ratio (Rr) was calculated by the following equation:  $Rr = W_f / W_0$ , where  $W_f$  and  $W_0$  are the weights of the sea cucumber before and after rehydration, respectively. Also, the patterns of change in weight ( $\Delta W$ ) and length ( $\Delta L$ ) of the individuals along the time period of rehydration (96 h) were evaluated.

## Results and discussion

The second methodology of rehydration (B) was more effective than the first one (A) taking into account the Rr, and length-weight recovering rates on *H. arguinensis* (Table 1): an average Rr of 3.775, was registered; this value being higher than the one obtained for *H. scabra* (Rr = 3.05) and *H. arguinensis* under rehydration protocol-A (Rr = 2.9). The highest value (Rr = 3.78) was relatively close to the Rr registered for *A. japonicus* (Rr = 4.07), which was obtained from a traditional rehydrated protocol in high-quality individuals (Zhang et al., 2016). It is important to highlight that the methodology used to produce beche-de-mer has an influence on its porous structure and therefore on its water holding capacity, which is linked to the rehydration ratio (Chong et al. 2015).





Figure 1. Rehydration protocols used on dried sea cucumbers (*Holothuria arguinensis* and *Holothuria scabra*).

**Table 1.** Values of length and weight increases ( $\Delta L$  and  $\Delta W$ ) and rehydration rate (Rr) for *Holothuria arguinensis* and *Holothuria scabra* specimens. Methods 'A' and 'B' are described in Material and methods section.

Individual	Species	Method	$\Delta L$ (%)	$\Delta W$ (%)	Rr
Individual A	<i>H. arguinensis</i>	A	28.00	65.57	2.90
Individual D	<i>H. arguinensis</i>	A	32.79	66.67	3.00
Individual B	<i>H. arguinensis</i>	B	37.04	73.49	3.77
Individual C	<i>H. arguinensis</i>	B	35.53	73.47	3.78
Individual E	<i>H. scabra</i>	A	34.92	67.21	3.05

**Figure 2.** Length (a) and weight (b) changes along rehydration protocol-A on individuals A, D, E (*Holothuria arguinensis* and *Holothuria scabra*); Length (c) and weight (d) changes along rehydration protocol-B on individuals B and C (*Holothuria arguinensis*).

*H. arguinensis*, which was rehydrated according the second methodology (B) showed average recovery rates of weight and length close to 73.4% and 36.3% respectively (Table 1; Figure 2). These values were also higher than rates registered with *H. scabra* (67.2% and 34.9%) or *H. arguinensis* rehydrated using methodology A without boiling (average values: 66.1% and 30.4%) (Table 1; Figure 2).

These results are very valuable to improve the rehydration protocols used on *Holothuria arguinensis*, as they allow for the maximum recovery rates of weight and length from the dried product.

## Acknowledgements

This research was supported by CUMARSUR (PTDC/MAR-BIO/5948/2014) and HOLREMAR projects funded by Fundação para a Ciência e a Tecnologia and Sayanes Mar S.L. Dr Mercedes González-Wangüemert was supported by FCT Investigator Programme-Career Development Contract (IF/00998/2014). J. Domínguez-Godino was supported by research fellow (CCMAR/BI/0007/2015).

## References

- Aydin M. 2008. The commercial sea cucumber fishery in Turkey. SPC Beche-de-mer Information Bulletin 28:40–41.
- Borrero-Pérez G., Gómez-Zurita J., González-Wangüemert M., Marcos C. and Pérez-Ruzafa A. 2010. Molecular systematic of the genus *Holothuria* in the Mediterranean and North-eastern Atlantic and a molecular clock for the diversification of the Holothuriidae (Echinodermata: Holothuroidea). Molecular Phylogenetics and Evolution 57:899–906.
- Chong N., Pindi W., Chye F.-Y., Shaarani S.M. and Lee J.S. 2015. Effects of drying methods on the quality of dried sea cucumbers from Sabah – A review. International Journal of Novel Research in Life Sciences 2:49–64.
- Duan X., Zhang M., Li X.L. and Mujumdar A.S. 2008. Microwave freeze drying of sea cucumber coated with nanoscale silver. Drying Technology 26(4):413–19.
- Fukunaga T., Matsumoto M., Murakami T. and Hatae K. 2004. Effects of soaking conditions on the texture of dried sea cucumber. Fishery Science 70(2):319–325.
- Geng S., Wang H., Wang X., Ma X., Xiao S., Wang J. and Ta M. 2015. A non-invasive NMR and MRI method to analyze the rehydration of dried sea cucumber. Analytical Methods, DOI 10.1039/c4ay03007a
- González-Wangüemert M. and Borrero-Pérez G. 2012. A new record of *Holothuria arguinensis* colonizing the Mediterranean Sea. Marine Biodiversity Records 5: e105.
- González-Wangüemert M., Domínguez-Godino J. and Cánovas F. 2018. The fast development of sea cucumber fisheries in the Mediterranean and NE Atlantic waters: from a new marine resource to its over-exploitation. Ocean and Coastal Management Journal 151:165–77.
- González-Wangüemert M., Valente S., Henriques F., Domínguez-Godino J. and Serrão E. 2016. Setting preliminary biometric baselines for new target sea cucumbers species of the NE Atlantic and Mediterranean fisheries. Fisheries Research 179:57–66.
- Hong J., Li D., Zhang H. and Chen L. 2014. Effect on quality of dried sea cucumber with different rehydration conditions. Journal of Fujian Fisheries 36(1):29–34.
- Liu H. and Ko W. 2002. Compositional analyses of dried sea cucumber *Metriatyla scabra* after rehydration. Turkish Journal of Agriculture and Forestry 51(4):1–8.
- Mezali K. and Thandar AS. 2014. First record of *Holothuria (Roweothuria) arguinensis* (Echinodermata: Holothuroidea: Aspidochirotida: Holothuriidae) from the Algerian coastal waters. Marine Biodiversity Records 7: e40.
- Micael J., Alves M.J., Jones M.B. and Costa A.C. 2012. Diversity of shallow-water asteroids (Echinodermata) in the Azorean Archipelago Research Articles. Marine Biodiversity Records 5: doi:10.17/S1755267211000534.
- Pereira M. 1997. Checklist of the littoral echinoderms of the Azores. Açoreana 8:331–37.
- Purcell S.W., Choo P.S., Akamine J., Fabinyi M. 2014. Alternative product forms, consumer packaging and extracted derivatives of tropical sea cucumbers. SPC Beche-de-mer Information Bulletin 34:47–52.
- Purcell S.W., Mercier A., Conand C., Hamel J.F., Toral-Granda V., Lovatelli A. and Uthicke S. 2013. Sea cucumber fisheries: global analysis of stocks, management measures and drivers of overfishing. Fish and Fisheries 14:34–59.
- Purcell S.W., Samyn Y. and Conand C. 2012. Commercially important sea cucumbers of the World. FAO, Rome. FAO Species Catalogue for Fishery Purposes No. 6. 190 p.
- Roggatz C., González-Wangüemert M., Pereira H., Rodrigues M., da Silva M., Barreira L., Varela J. and Custodio L. 2016. First report of the nutritional profile and antioxidant potential of *Holothuria arguinensis*, a new resource for aquaculture in Europe. Natural Product Research Journal 30:2034–40.
- Sze C.P. and Conand C. 2015. Sea cucumber products at the China Fisheries Seafood Expo 2014. SPC Beche-de-mer Information Bulletin 35:60–63.
- Xiang Y., Su X., Dong M. and Chen C. 2007. Study on water immersion technology of salted sea cucumber. Food Science 28(12):153–56.
- Zhang L., Huang X., Miao S., Zeng S., Zhang Y. and Zheng B. 2016. Influence of ultrasound on the rehydration of dried sea cucumber (*Stichopus japonicus*). Journal of Food Engineering: <http://dx.doi.org/10.1016/j.jfoodeng.2016.01.024>