

Public Health

- [Life expectancy at birth \(Years\)](#)
- [Life expectancy at age 40 \(e/40\) \(Years\)](#)
- [Prevalence of Adult Diabetes \(age 25-64\)](#)
- [Proportion of 1 year old children immunized against measles \(%\)](#)
- [Contraceptive prevalence rate \(%\)](#)
- [Antenatal care coverage \(% of women who had at least one antenatal session with a skilled provider\)](#)
- [Low birth weight babies \(%\)](#)
- [Prevalence of overweight or obese adults - BMI \$\geq\$ 25.0](#)
- [Prevalence of obese children - 0-5 year olds \(+2SD from median for BMI by height\) \(%\)](#)
- [Proportion of Households with access to improved drinking water \(%\)](#)
- [Proportion of Households with access to improved sanitation \(%\)](#)
- [National Health Accounts - Government expenditure on health as % of total government expenditure](#)
- [National Health Accounts - Per capita total expenditure on health \(US\\$\)](#)
- [Sustainability of national health systems \(prop of annual health expenditure from international aid\) \(%\)](#)
- [Proportion of one year olds fully immunized against EPI target diseases \(3 doses diptheria, tetanus, toxoid & pertussis\) \(%\)](#)
- [Proportion of one year olds fully immunized against EPI target diseases \(3 doses hepatitis B - HepB3\) \(%\)](#)
- [Prevalence of obese adults \(aged 25-64\) BMI \$\geq\$ 30](#)
- [Maternal Mortality \(Number of Deaths\)](#)
- [Prevalence of obese children - 13-15 year olds \(+2SD from median for BMI by age\) \(%\)](#)
- [Total Fertility Rate](#)
- Maternal mortality ratio
- Proportion of births attended
- Under-five mortality rate
- Neonatal mortality rate
- Tuberculosis incidence per 100,000 population
- Malaria incidence per 1,000 population
- Number of people requiring interventions against neglected tropical diseases
- Mortality rate attributed to cardiovascular disease, cancer, diabetes or chronic respiratory disease
- Harmful use of alcohol, defined according to the national context as alcohol per capita consumption (aged 15 years and older) within a calendar year in litres of pure alcohol
- Proportion of women of reproductive age (aged 15-49 years) who have their need for family planning satisfied with modern methods
- Adolescent birth rate (aged 10-14 years; aged 15-19 years) per 1,000 women in that age group

- Coverage of essential health services (defined as the average coverage of essential services based on tracer interventions that include reproductive, maternal, newborn and child health, infectious diseases, non-communicable diseases and service capacity and access, among the general and the most disadvantaged population)
- Mortality rate attributed to unsafe water, unsafe sanitation and lack of hygiene (exposure to unsafe Water, Sanitation and Hygiene for All (WASH) services)
- Age-standardized prevalence of current tobacco use among persons aged 15 years and older
- Health worker density and distribution
- International Health Regulations (IHR) capacity and health emergency preparedness