


B E C H E - D E - M E R

NEWS

*Bohadschia argus*

The dive fishery of sea-cucumbers in Washington State

by A. Bradbury¹ and C. Conand²¹Washington Department of Fisheries, USA²Labo. Océanographie Biologie, UBO, France

The landings of *Parastichopus californicus* from a commercial dive fishery in Washington State have been recorded since 1971.

This fishery is particularly noteworthy as the five longitudinal muscles are stripped from the body wall, frozen and exported mainly to Asian markets (Taiwan). The body wall itself is also processed into beche-de-mer for export.

In a first review, Sloan (1985, 1986) states that 'the fishery is still characterized by incomplete market development which imparts instability and causes non stock-related fluctuations in landings'. Until 1987, there were no seasonal nor areal restrictions for the divers. The development of the fishery was assessed through mandatory log books filled in by divers holding a licence.

Following signs of overfishing, the Washington State Department of Fisheries divided state waters into four harvest areas - San Juan Islands (1), Strait of Juan de Fuca (2), Central Puget Sound (3) and South Puget Sound/Hood Canal (4). Seasons now rotate, so each area goes unfished for 3.5 years after a six-month harvest. A seasonal restriction has also been implemented, with a six month fishing season extending from May to October.

The closed period was based on the seasonal atrophy of the visceral organs as described by Fankboner and Cameron (1985). These rotating area closures give sea-cucumbers the chance to spawn at least once before being harvested. The recent progress in understanding the biology of this species (Cameron and

Fankboner, 1989) has permitted a more scientific basis for the management of this fishery. Limited-entry legislation has also been approved in 1990 and the number of boats licensed is now declining.

The annual landings of the Washington sea cucumber fishery are shown in Figure 1 though these figures should be viewed with caution, due to unreporting. Landings stayed under 10 tons until 1977, then fluctuated for ten years at an intermediate level and not exceeding 200 tons. The figure shows a large increase in landings between 1988 and 1990, with the record catch, in 1989, of over 1,000 tons.

A summary of the production from the beche-de-mer fishery is given in Table 1. The value per kilogram has more than doubled during the last four years, making the fishery attractive for divers. Since 1983, the number of boats, fishing effort (dive




Figure 1. Evolution of *Parastichopus* landings from the Pacific coast of North America.

hours) and commercial buyers have all increased markedly. The CPUE, in number of sea cucumbers, although variable, is decreasing in the different districts forcing the divers to fish in deeper waters.

Among the different measures taken to prevent overfishing, the limited-entry legislation is probably the most important conservation strategy.

Table 1. Sea cucumber exploitation by divers in Washington State (U.S.A).

Year	Catch (kg)	Value(1)		Number boats (2)	District 1			District 2			District 3			District 4		
		Total (US\$ *10^6)	per kg (US\$)		Effort (3)	CPUE (4) n	kg	Effort (3)	CPUE (4) n	kg	Effort (3)	CPUE (4) n	kg	Effort (3)	CPUE (4) n	kg
1971	3,629															
1972	2,722															
1973	4,536															
1974	-															
1975	1,361															
1976	6,804															
1977	28,577															
1978	57,607															
1979	107,050															
1980	190,966															
1981	125,194															
1982	12,247															
1983	170,554			9				99	530		281	275				
1984	39,917			14	68	271		225	495		310	220		28	502	
1985	146,966			15	74	232		162	397		1,763	272		336	636	
1986	181,894			16	345	200		253	425		1,116	249		237	321	
1987(5)	127,008	0.89	0.7	25	837	248	69									
1988	871,366	1.07	1.23	78				6,492	381	106.1						
1989	1,146,701	1.87	1.63	125							11,437	239	66.7			
1990	992,794	1.75	1.76	79										5,833	264	73.5

- (1) Estimated from the average price paid per bucket of slit, drained cucumber
- (2) Number of boats making at least one landing
- (3) Total dive hours
- (4) Mean number (n) or weight (kg) per dive hour, from harvest logs
- (5) Since 1987, area openings are rotating (districts 1 to 4).

In the meantime, assessments of the present resources and of the holothurian population after the fishing period, are being conducted by fishery biologists.

At present, the Washington dive fishery is probably the best documented and most carefully managed sea-cucumber fishery.

References

Bradbury A., 1990. Management of the commercial dive fisheries for sea urchins and sea cucumbers. *In: Proceedings from the Forum on Puget Sound's Biological Resources. Status and Management.* Armstrong and Copping (eds), U.S. EPA Publication 910/9-90-001: 56-65.

Cameron J.L. & Fankboner P.V., 1989. Reproductive biology of the commercial sea cucumber

Parastichopus californicus (Echinodermata: Holothuroidea). II. Observations on the ecology of development, recruitment and juvenile life stage. *Journ. Exp. Mar. Biol. Ecol.*, 127: 43-67.

Fankboner P.V. & Cameron J.L., 1985. Seasonal atrophy of the visceral organs in a sea cucumber. *Can. J. Zool.*, 63: 2888-2892.

Sloan N.A., 1985. Echinoderm fisheries of the world: a review. *In: Echinodermata.* B. Keegan and B. O'Connor. Balkema, Rotterdam: 109-124.

Sloan N.A., 1986. World Jellyfish and Tunicate Fisheries and the Northeast Pacific Echinoderm Fishery. *In: G.S. Jamieson and N. Bourne (ed.). North Pacific Workshop on Stock Assessment and Management of Invertebrates.* *Can. Spec. Publ. Fish. Aquat. Sci.*, 92: 430p.

Measurement of size in live sea cucumbers

by M. A. Sewell,
Zoology Department,
University of Alberta, USA

The need to devise a suitable and reliable measure of size in live sea cucumbers has been a consistent problem for researchers investigating the population ecology of this group. This was noted by Alex Bradbury (Washington State Department of Fisheries) in a recent issue of *Beche-de-mer Information Bulletin*. Any person faced with a sea cucumber deflating in one's hands will probably despair of ever getting a repeatable measure of size.

Measurement has usually been based on two variables — total length and total weight. However, there are problems with both these measures. Length is difficult to measure because of the morphology and elasticity of the body wall. Variation is also observed in total weight due to differences in the amount of food in the gut, seasonal changes in the size of the alimentary canal and respiratory tree (in some species of sea cucumber), and the quantity of water contained within the respiratory tree (Choe, 1963; Conand, 1981, 1989).

For research on the aspidochirote *Stichopus mollis*, a reliable measure of size was needed for calculation of gonad indices and size-frequency distributions. The method devised required that the animals be brought to the laboratory, but provided fairly reliable results. Although this method is described in a recent paper (Sewell, 1990), details of the methods are given here.

To get a reliable measure of size, the animals were placed in aquaria for 24 hours in order to clear the sediment from the alimentary canal. Each animal was put in a small mesh bag constructed from surplus onion sacks (mesh size 5 mm). The bags were approximately 20 x 20 cm square, just large enough to contain the animals. In the area studied, sexually mature *Stichopus mollis* were 13–24 centimetres in length. A large metal frame (a salvaged rectangular quadrat) was hung over a pool of running sea water (500 litres) as shown in Figure 1.

Then 'hangers' were constructed from two squeeze-type clothes pegs connected by string. The bag was closed by this hanger, and suspended from the metal frame so that the animal was totally immersed in the running sea water. Animals suspended in this fashion shed faeces to the bottom of the tank, and could not reconsume the sediments. After 24 hours, each animal was transferred to a small laboratory aquarium (15 l) of aerated sea water for 15 minutes. This allowed the animal to achieve a natural body form after being held in the mesh




Figure 1. Side-view of aquarium showing position of holding bags. (Not to scale)

bags. Most animals started respiratory movements soon after entering the tank, and began moving tentacles after 5–10 minutes.

Two measures of size were made — total length and weight. Each sea cucumber was removed from the aquarium and its length measured dorsally from the anus to the centre of the tentacular crown with a dressmaker's tape to the nearest 0.5 cm. The animal was stimulated to expel the excess water remaining in the respiratory trees by gently squeezing the posterior half, and then quickly blotted dry and weighed.

To obtain an estimate of the errors associated with the total length and total weight measures used, ten sea cucumbers were measured ten times at 15-minute intervals. Details of the analysis and table of results are included in Sewell (1990). Raw data are included in Table 1.

The results showed that total length was not a very reliable measure of size (Sewell, 1990). Individual sea cucumbers varied up to 6 cm in length. In this study, however, there was no correlation between the size of animal measured in the field prior to collection (i.e. underwater), and the mean of the ten repeated measures in the laboratory. I, therefore, do not recommend the use of length to measure size.

Total weight appeared to be a reliable measure of size in *Stichopus mollis*, provided that sediments were removed from the gut and excess water squeezed from the respiratory trees. Analysis showed that a single measure had a reliability of approximately 95 per cent in estimating the mean

Table 1: Repeated measures analysis of total weight and total length in *Stichopus mollis*

Animal	Weight range (g)	Mean (g)	S.D. (g)	Length range (cm)	Mean (cm)	S.D. (cm)
1	178.1 - 205.0	189.4	8.3	17.5 - 20.5	18.5	1.0
2	178.2 - 206.1	198.3	8.8	17.0 - 21.5	19.4	1.3
3	138.6 - 156.9	145.6	6.1	16.5 - 19.5	17.9	0.9
4	146.4 - 159.0	154.7	6.2	14.5 - 20.0	16.9	1.9
5	173.6 - 194.4	183.9	7.1	16.0 - 22.0	18.1	1.9
6	183.2 - 193.1	188.1	3.4	18.0 - 19.5	18.9	0.5
7	103.8 - 115.6	107.1	3.4	12.5 - 17.0	14.8	1.4
8	148.6 - 166.1	156.8	5.7	16.0 - 19.5	18.1	0.9
9	154.7 - 163.8	158.7	3.1	15.5 - 19.0	17.7	1.1
10	175.9 - 203.7	191.6	7.5	18.5 - 20.0	19.4	0.5

of the ten measures (Sewell, 1990). A high correlation was also found between total weight and the drained weight used in the calculation of the gonad index ($r=0.95$, $n=480$), suggesting that total weight was a reasonably accurate and non-destructive measure of size in this species.

So in conclusion, a reasonable measure of weight could be obtained by the following protocol for *Stichopus mollis*

- (i) Empty the alimentary canal in such a way that sea cucumbers cannot reconsume the sediments released as faeces;
- (ii) Allow a relaxation period in the tank;
- (iii) Squeeze excess water from the respiratory trees;
- (iv) Blot quickly and weigh.

These methods were found to be suitable for calculation of gonad indices; however, if animals are to be reweighed for estimates of growth a number of measurements should be taken, and a mean calculated. The method described here reduces some of the variation, by at least ensuring that all the animals have empty guts before measurement.

References:

- Choe, S.(1963). *Namako no kenkyu* (Biology of the Japanese common sea cucumber *Stichopus japonicus*, Selenka). Pusan National University, Japan, 226 pp. (in Japanese with English summary).
- Conand, C. 1989. Les holothuries Aspidochirotes du lagon de Nouvelle-Calédonie: Biologie, écologie et exploitation, Etudes et thèses, ORSTROM., Paris, 393 pp.
- Conand, C. 1981. Sexual cycle of three commercially important holothurian species (Echinodermata) from the lagoon of New Caledonia, *Bull. Mar. Sci.*, 31 (1981) 523-543.
- Sewell, M.A. (1990). Aspects of the ecology of *Stichopus mollis* (Echinodermata: Holothuroidea) in north-eastern New Zealand. *N.Z. J. Mar. F.W. Res.* 24: 97-103.

Recent trends in sea cucumbers exploitation in New Caledonia

by C. Conand¹ and C. Hoffschir²

¹Labo. Océanographie Biologie, UBO, France
²O.R.S.T.O.M, Nouméa, Nouvelle Calédonie

The history of the sea cucumber fishery in New Caledonia has been traced back to the nineteenth century from various documents and statistics. The exports have shown wide fluctuations in connection with political and socio-economic events, as well as with the status of the biological resource (Conand, 1989, 1990).

Evolution of the exports

The most recent period of revival started in 1983, when a few New Caledonians of Chinese origin undertook to organise the fishing, processing and trade. As the catches in

New Caledonia are all exported, the export statistics provide easily obtainable accurate data.

Table 1 shows the fluctuations in tonnage between 1983 and 1990. For this period, the mean annual export was 103 tons. Production was not constant but oscillated between years, giving peaks in landed volume during 1984, 1986, 1988 and 1990.

Table 1. Evolution of the recent beche-de-mer exports from New Caledonia

Year	1983	1984	1985	1986	1987	1988	1989	1990
Tonnage	15	150	89	180	77	135	55	126
exporters	2	4	7	6	4	3	3	4